

# A HILBERT-TYPE INTEGRAL INEQUALITY IN THE WHOLE PLANE RELATED TO THE HYPERGEOMETRIC FUNCTION AND THE BETA FUNCTION

Michael Th. Rassias<sup>1\*</sup> and Bicheng Yang<sup>2</sup>

<sup>1\*</sup>. Department of Mathematics, Princeton University,  
Fine Hall, Washington Road, Princeton, NJ 08544-1000, USA  
& Department of Mathematics, ETH-Zürich,  
Rämistrasse 101, 8092 Zürich, Switzerland,  
michailrassias@math.princeton.edu michail.rassias@math.ethz.ch

<sup>2</sup>. Department of Mathematics, Guangdong University of  
Education, Guangzhou, Guangdong 510303, P. R. China  
E-mail: <sup>1\*</sup>. Corresponding author: michael.rassias@math.ethz.ch  
2. bcyang@gdei.edu.cn bcyang818@163.com

## Abstract

A new Hilbert-type integral inequality in the whole plane with the non-homogeneous kernel and parameters is given. The constant factor related to the hypergeometric function and the beta function is proved to be the best possible. As applications, equivalent forms, the reverses, some particular examples, two kinds of Hardy-type inequalities, and operator expressions are considered.

**Key words:** Hilbert-type integral inequality; weight function; equivalent form; hypergeometric function; beta function;

**2000 Mathematics Subject Classification.** 26D15 33B15

## 1 Introduction

If  $f(x), g(y) \geq 0$ , satisfy

$$0 < \int_0^\infty f^2(x)dx < \infty$$

and

$$0 < \int_0^\infty g^2(y)dy < \infty,$$

---

<sup>0</sup>**Foundation item:** This work is supported by the National Natural Science Foundation of China (No. 61370186), and 2013 Knowledge Construction Special Foundation Item of Guangdong Institution of Higher Learning College and University (No. 2013KJCX0140).

then we have the following Hilbert's integral inequality (cf. [1]):

$$\int_0^\infty \int_0^\infty \frac{f(x)g(y)}{x+y} dx dy < \pi \left( \int_0^\infty f^2(x) dx \int_0^\infty g^2(y) dy \right)^{\frac{1}{2}}, \quad (1.1)$$

where the constant factor  $\pi$  is the best possible. The inequality (1.1) is very important in Mathematical Analysis and its applications (cf. [1], [2]). In recent years, by the use of the method of weight functions, a number of extensions of (1.1) were given by Yang (cf. [3]). Noticing that inequality (1.1) is a homogenous kernel of degree -1, in 2009, a survey of the study of Hilbert-type inequalities with the homogeneous kernels of degree equal to negative numbers and some parameters is given in [4]. Recently, some inequalities with the homogenous kernels of degree 0 and non-homogenous kernels have been proved (cf. [5]-[10]). Other kinds of Hilbert-type inequalities are shown in [11]-[16]. All of the above integral inequalities are constructed in the quarter plane of the first quadrant.

In 2007, Yang [17] presented a new Hilbert-type integral inequality in the whole plane, as follows:

$$\begin{aligned} & \int_{-\infty}^\infty \int_{-\infty}^\infty \frac{f(x)g(y)}{(1+e^{x+y})^\lambda} dx dy \\ & < B\left(\frac{\lambda}{2}, \frac{\lambda}{2}\right) \left( \int_{-\infty}^\infty e^{-\lambda x} f^2(x) dx \int_{-\infty}^\infty e^{-\lambda y} g^2(y) dy \right)^{\frac{1}{2}}, \end{aligned} \quad (1.2)$$

where the constant factor  $B(\frac{\lambda}{2}, \frac{\lambda}{2})(\lambda > 0)$  is the best possible.

If  $0 < \lambda < 1$ ,  $p > 1$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ , Yang [20] derived another new Hilbert-type integral inequality in the whole plane. Namely, he proved that

$$\begin{aligned} & \int_{-\infty}^\infty \int_{-\infty}^\infty \frac{1}{|1+xy|^\lambda} f(x)g(y) dx dy \\ & < k_\lambda \left[ \int_{-\infty}^\infty |x|^{p(1-\frac{\lambda}{2})-1} f^p(x) dx \right]^{\frac{1}{p}} \left[ \int_{-\infty}^\infty |y|^{q(1-\frac{\lambda}{2})-1} g^q(y) dy \right]^{\frac{1}{q}}, \end{aligned} \quad (1.3)$$

where the constant factor

$$k_\lambda = B\left(\frac{\lambda}{2}, \frac{\lambda}{2}\right) + 2B\left(1-\lambda, \frac{\lambda}{2}\right)$$

is still the best possible. Furthermore, Yang et al. [19]-[28] proved as well some new Hilbert-type integral inequalities in the whole plane.

In this paper, using methods from Real Analysis and by estimating the weight functions, a new Hilbert-type integral inequality in the whole plane with the non-homogeneous kernel and multi-parameters is shown, which gives an extension of (1.3). The constant factor related to the hypergeometric function and the beta function is proved to be the best possible. As applications, equivalent forms, the reverses, some particular examples, two kinds of Hardy-type inequalities, and operator expressions are also considered.

## 2 Some Lemmas

Initially, we introduce the following formula of the hypergeometric function  $F$  (cf. [29]):  
If  $\operatorname{Re}(\gamma) > \operatorname{Re}(\theta) > 0$ ,  $|\arg(1-z)| < \pi$ ,  $(1-zt)^{-\alpha}|_{z=0} = 1$ , then

$$F(\alpha, \theta, \gamma, z) := \frac{\Gamma(\gamma)}{\Gamma(\theta)\Gamma(\gamma-\theta)} \int_0^1 t^{\theta-1} (1-t)^{\gamma-\theta-1} (1-zt)^{-\alpha} dt,$$

where,

$$\Gamma(\eta) = \int_0^\infty x^{\eta-1} e^{-x} dx (\operatorname{Re}(\eta) > 0)$$

is the gamma function. In particular, for  $z = -1$ ,  $\gamma = \theta + 1$  ( $\theta > 0$ ),  $\alpha \in \mathbf{R}$ , we have

$$\int_0^1 t^{\theta-1} (1+t)^{-\alpha} dt = \frac{1}{\theta} F(\alpha, \theta, 1+\theta, -1) \in \mathbf{R}_+. \quad (2.1)$$

**Lemma 2.1.** *If  $\beta > -1$ ,  $\mu, \sigma > -\beta$ ,  $\mu + \sigma = \lambda < 1 - \beta$ ,  $\delta \in \{-1, 1\}$ , we define two weight functions  $\omega(\sigma, y)$  and  $\varpi(\sigma, x)$  as follows:*

$$\omega(\sigma, y) : = \int_{-\infty}^\infty \frac{(\min\{1, |x^\delta y|\})^\beta}{|1+x^\delta y|^{\lambda+\beta}} \frac{|y|^\sigma}{|x|^{1-\delta\sigma}} dx (y \in \mathbf{R} \setminus \{0\}), \quad (2.2)$$

$$\varpi(\sigma, x) : = \int_{-\infty}^\infty \frac{(\min\{1, |x^\delta y|\})^\beta}{|1+x^\delta y|^{\lambda+\beta}} \frac{|x|^{\delta\sigma}}{|y|^{1-\sigma}} dy (x \in \mathbf{R} \setminus \{0\}). \quad (2.3)$$

Then we have

$$\begin{aligned} \omega(\sigma, y) &= \varpi(\sigma, x) = K(\sigma) := \frac{1}{\beta + \sigma} F(\lambda + \beta, \beta + \sigma, 1 + \beta + \sigma, -1) \\ &\quad + \frac{1}{\beta + \mu} F(\lambda + \beta, \beta + \mu, 1 + \beta + \mu, -1) \\ &\quad + B(1 - \lambda - \beta, \beta + \sigma) + B(1 - \lambda - \beta, \beta + \mu) \in \mathbf{R}_+. \end{aligned} \quad (2.4)$$

*Proof.* (i) For  $\delta = 1$ , by (2.2) it follows that

$$\omega(\sigma, y) = \int_{-\infty}^\infty \frac{(\min\{1, |xy|\})^\beta}{|1+xy|^{\lambda+\beta}} \frac{|y|^\sigma}{|x|^{1-\sigma}} dx.$$

(a) If  $y < 0$ , setting  $u = xy$ , we obtain

$$\begin{aligned} \omega(\sigma, y) &= \int_{-\infty}^\infty \frac{(\min\{1, |u|\})^\beta}{|1+u|^{\lambda+\beta}} \frac{(-y)^\sigma}{|u/y|^{1-\sigma}} \frac{1}{y} du \\ &= \int_{-\infty}^\infty \frac{(\min\{1, |u|\})^\beta}{|1+u|^{\lambda+\beta}} \frac{(-y)^\sigma (-y)^{1-\sigma}}{|u|^{1-\sigma}} \frac{1}{(-y)} du \\ &= \int_{-\infty}^\infty \frac{(\min\{1, |u|\})^\beta |u|^{\sigma-1}}{|1+u|^{\lambda+\beta}} du; \end{aligned}$$

(b) if  $y > 0$ , setting  $u = xy$ , it yields

$$\omega(\sigma, y) = \int_{-\infty}^\infty \frac{(\min\{1, |u|\})^\beta}{|1+u|^{\lambda+\beta}} \frac{y^\sigma du}{|u/y|^{1-\sigma} y} = \int_{-\infty}^\infty \frac{(\min\{1, |u|\})^\beta |u|^{\sigma-1}}{|1+u|^{\lambda+\beta}} du.$$

(ii) For  $\delta = -1$ , setting  $X = x^{-1}$ , we obtain

$$\begin{aligned}
\omega(\sigma, y) &= \int_{-\infty}^0 \frac{(\min\{1, |x^{-1}y|\})^\beta}{|1+x^{-1}y|^{\lambda+\beta}} \frac{|y|^\sigma}{|x|^{1+\sigma}} dx + \int_0^\infty \frac{(\min\{1, |x^{-1}y|\})^\beta}{|1+x^{-1}y|^{\lambda+\beta}} \frac{|y|^\sigma}{|x|^{1+\sigma}} dx \\
&= \int_0^{-\infty} \frac{(\min\{1, |Xy|\})^\beta}{|1+Xy|^{\lambda+\beta}} \frac{-|y|^\sigma dX}{|X^{-1}|^{1+\sigma} X^2} + \int_\infty^0 \frac{(\min\{1, |Xy|\})^\beta}{|1+Xy|^{\lambda+\beta}} \frac{-|y|^\sigma dX}{|X^{-1}|^{1+\sigma} X^2} \\
&= \int_{-\infty}^0 \frac{(\min\{1, |Xy|\})^\beta}{|1+Xy|^{\lambda+\beta}} \frac{|y|^\sigma dX}{|X|^{1-\sigma}} + \int_0^\infty \frac{(\min\{1, |Xy|\})^\beta}{|1+Xy|^{\lambda+\beta}} \frac{|y|^\sigma dX}{|X|^{1-\sigma}} \\
&= \int_{-\infty}^\infty \frac{(\min\{1, |xy|\})^\beta}{|1+xy|^{\lambda+\beta}} \frac{|y|^\sigma}{|x|^{1-\sigma}} dx.
\end{aligned}$$

Hence, for  $\delta \in \{-1, 1\}$ , we obtain the following expression:

$$\begin{aligned}
\omega(\sigma, y) &= \int_{-\infty}^\infty \frac{(\min\{1, |u|\})^\beta |u|^{\sigma-1}}{|1+u|^{\lambda+\beta}} du = K_1(\sigma) + K_2(\sigma), \\
K_1(\sigma) &:= \int_{-1}^1 \frac{(\min\{1, |u|\})^\beta |u|^{\sigma-1}}{(1+u)^{\lambda+\beta}} du, \\
K_2(\sigma) &:= \int_{\mathbf{R} \setminus (-1, 1)} \frac{(\min\{1, |u|\})^\beta |u|^{\sigma-1}}{|1+u|^{\lambda+\beta}} du.
\end{aligned}$$

In view of (2.1), and the following formula of the beta function (cf. [29]):

$$B(p, q) := \int_0^1 \frac{v^{p-1}}{(1-v)^{1-q}} dv \quad (p, q > 0),$$

we obtain

$$\begin{aligned}
K_1(\sigma) &= \int_{-1}^0 \frac{(-u)^{\beta+\sigma-1}}{(1+u)^{\lambda+\beta}} du + \int_0^1 \frac{u^{\beta+\sigma-1}}{(1+u)^{\lambda+\beta}} du \\
&= \int_0^1 \frac{v^{\beta+\sigma-1}}{(1-v)^{\lambda+\beta}} dv + \int_0^1 \frac{u^{\beta+\sigma-1}}{(1+u)^{\lambda+\beta}} du \\
&= B(1-\lambda-\beta, \beta+\sigma) + \frac{1}{\beta+\sigma} F(\lambda+\beta, \beta+\sigma, 1+\beta+\sigma, -1). \quad (2.5)
\end{aligned}$$

Setting  $v = \frac{1}{u}$ , it follows that

$$\begin{aligned}
K_2(\sigma) &= \int_{-\infty}^{-1} \frac{(\min\{1, (-u)\})^\beta (-u)^{\sigma-1}}{|1+u|^{\lambda+\beta}} du + \int_1^\infty \frac{(\min\{1, u\})^\beta u^{\sigma-1}}{|1+u|^{\lambda+\beta}} du \\
&= - \int_0^{-1} \frac{(\min\{1, (-\frac{1}{v})\})^\beta (-\frac{1}{v})^{\sigma-1}}{|1+\frac{1}{v}|^{\lambda+\beta} v^2} dv - \int_1^0 \frac{(\min\{1, \frac{1}{v}\})^\beta (\frac{1}{v})^{\sigma-1}}{|1+\frac{1}{v}|^{\lambda+\beta} v^2} dv \\
&= \int_{-1}^0 \frac{(\min\{1, (-v)\})^\beta (-v)^{\lambda-\sigma-1}}{|1+v|^{\lambda+\beta}} dv + \int_0^1 \frac{(\min\{1, v\})^\beta v^{\lambda-\sigma-1}}{|1+v|^{\lambda+\beta}} dv \\
&= \int_{-1}^1 \frac{(\min\{1, |v|\})^\beta |v|^{\mu-1}}{(1+v)^{\lambda+\beta}} dv = K_1(\mu) \\
&= B(1-\lambda-\beta, \beta+\mu) + \frac{1}{\mu} F(\lambda+\beta, \beta+\mu, 1+\beta+\mu, -1). \quad (2.6)
\end{aligned}$$

Setting  $u = x^\delta y$  in (2.3), for  $x < 0$  ( $x > 0$ ), we also get

$$\varpi(\sigma, x) = \int_{-\infty}^{\infty} \frac{(\min\{1, |u|\})^\beta |u|^{\sigma-1}}{|1+u|^{\lambda+\beta}} du = K(\sigma).$$

Hence we have (2.4). □

*Remark 2.2.* By Taylor's formula, we obtain

$$\begin{aligned} & \frac{1}{\beta + \sigma} F(\lambda + \beta, \beta + \sigma, 1 + \beta + \sigma, -1) \\ &= \int_0^1 \frac{u^{\beta+\sigma-1} du}{(1+u)^{\lambda+\beta}} = \int_0^1 \sum_{k=0}^{\infty} \binom{-\lambda-\beta}{k} u^{k+\beta+\sigma-1} du \\ &= \int_0^1 \sum_{k=0}^{\infty} (-1)^k \binom{\lambda+\beta+k-1}{k} u^{k+\beta+\sigma-1} du \\ &= \int_0^1 \sum_{k=0}^{\infty} \left[ \binom{\lambda+\beta+2k-1}{2k} - \binom{\lambda+\beta+2k}{2k+1} u \right] u^{2k+\beta+\sigma-1} du. \end{aligned}$$

Since we have

$$\binom{\lambda+\beta+2k-1}{2k} - \binom{\lambda+\beta+2k}{2k+1} u = \left[ 1 - \frac{(\lambda + \beta + 2k)u}{2k+1} \right] \binom{\lambda+\beta+2k-1}{2k},$$

there exists a large number  $k_0 \in \mathbf{N}_0 = \mathbf{N} \cup \{0\}$ , such that  $\lambda + \beta + 2k_0 > 0$ , and for any  $s \in \mathbf{N}$ ,

$$\begin{aligned} & \binom{\lambda+\beta+2(k_0+s)-1}{2(k_0+s)} - \binom{\lambda+\beta+2(k_0+s)+1}{2(k_0+s+1)} u \\ &= \left[ 1 - \frac{(\lambda + \beta + 2k_0 + 2s)u}{2(k_0 + s) + 1} \right] \binom{\lambda+\beta+2(k_0+s)-1}{2(k_0+s)} \\ &= \left[ 1 - \frac{(\lambda + \beta + 2k_0 + 2s)u}{2(k_0 + s) + 1} \right] \\ & \quad \times \frac{\lambda + \beta + 2k_0 + 2s - 1}{2k_0 + 2s} \dots \frac{\lambda + \beta + 2k_0}{2k_0 + 1} \binom{\lambda+\beta+2k_0-1}{2k_0}, \\ &= 1 - \frac{(\lambda + \beta + 2k_0 + 2s)u}{2(k_0 + s) + 1} \\ &\geq 1 - \frac{\lambda + \beta + 2k_0 + 2s}{2(k_0 + s) + 1} = \frac{1 - \lambda - \beta}{2(k_0 + s) + 1} > 0 (u \in (0, 1]). \end{aligned}$$

It follows that for any  $s \in \mathbf{N}$ , we have

$$\operatorname{sgn} \left( \binom{\lambda+\beta+2(k_0+s)-1}{2(k_0+s)} - \binom{\lambda+\beta+2(k_0+s)+1}{2(k_0+s+1)} u \right) = \operatorname{sgn} \left( \binom{\lambda+\beta+2k_0-1}{2k_0} \right).$$

By Lebesgue's term by term integration theorem (cf. [31]), we have

$$\begin{aligned} & \frac{1}{\sigma} F(\lambda + \beta, \beta + \sigma, 1 + \beta + \sigma, -1) \\ &= \sum_{k=0}^{\infty} \int_0^1 \left[ \binom{\lambda+\beta+2k-1}{2k} - \binom{\lambda+\beta+2k}{2k+1} u \right] u^{2k+\beta+\sigma-1} du \\ &= \sum_{k=0}^{\infty} \binom{-\lambda-\beta}{k} \int_0^1 u^{k+\beta+\sigma-1} du = \sum_{k=0}^{\infty} \frac{1}{k + \beta + \sigma} \binom{-\lambda-\beta}{k}. \end{aligned}$$

Similarly, since

$$\begin{aligned} (-1)^k \binom{-\lambda-\beta}{k} &= (-1)^k \frac{(-\lambda-\beta)(-\lambda-\beta-1)\cdots(-\lambda-\beta-k+1)}{k!} \\ &= (-1)^{2k} \frac{(\lambda+\beta)(\lambda+\beta+1)\cdots(\lambda+\beta+k-1)}{k!} = \binom{\lambda+\beta+k-1}{k}, \end{aligned}$$

we obtain

$$\begin{aligned} B(1-\lambda-\beta, \beta+\sigma) &= \int_0^1 \frac{u^{\beta+\sigma-1} du}{(1-u)^{\lambda+\beta}} = \int_0^1 \sum_{k=0}^{\infty} (-1)^k \binom{-\lambda-\beta}{k} u^{k+\beta+\sigma-1} du \\ &= \int_0^1 \sum_{k=0}^{\infty} \binom{\lambda+\beta+k-1}{k} u^{k+\beta+\sigma-1} du. \end{aligned}$$

There exists a large number  $k_1 \in \mathbf{N}$ , such that  $\lambda + \beta + k_1 > 0$ .

Hence, for any  $s \in \mathbf{N}$ , we have

$$\binom{\lambda+\beta+k_1+s-1}{k_1+s} = \frac{\lambda+\beta+k_1+s-1}{k_1+s} \cdots \frac{\lambda+\beta+k_1}{k_1+1} \binom{\lambda+\beta+k_1-1}{k_1},$$

and then it follows that

$$\operatorname{sgn} \binom{\lambda+\beta+k_1+s-1}{k_1+s} = \operatorname{sgn} \binom{\lambda+\beta+k_1-1}{k_1}.$$

Still by Lebesgue's term by term integration theorem, we obtain

$$B(1-\lambda-\beta, \beta+\sigma) = \sum_{k=0}^{\infty} \binom{\lambda+\beta+k-1}{k} \int_0^1 u^{k+\beta+\sigma-1} du = \sum_{k=0}^{\infty} \frac{(-1)^k}{k+\beta+\sigma} \binom{-\lambda-\beta}{k}.$$

Hence, we deduce the following series expressions:

$$K_1(\sigma) = 2 \sum_{k=0}^{\infty} \frac{1}{2k+\beta+\sigma} \binom{-\lambda-\beta}{2k}, \quad (2.7)$$

$$K_2(\sigma) = 2 \sum_{k=0}^{\infty} \frac{1}{2k+\beta+\mu} \binom{-\lambda-\beta}{2k}, \quad (2.8)$$

$$K(\sigma) = 2 \sum_{k=0}^{\infty} \frac{4k+2\beta+\lambda}{(2k+\beta+\sigma)(2k+\beta+\mu)} \binom{-\lambda-\beta}{2k}. \quad (2.9)$$

**Lemma 2.3.** Suppose that  $p > 1, \frac{1}{p} + \frac{1}{q} = 1, \beta > -1, \mu, \sigma > -\beta, \mu + \sigma = \lambda < 1 - \beta, \delta \in \{-1, 1\}, K(\sigma)$  as indicated by (2.4) (or (2.9)).

If  $f(x)$  is a non-negative measurable function in  $\mathbf{R}$ , then we have

$$\begin{aligned} J &: = \int_{-\infty}^{\infty} |y|^{p\sigma-1} \left[ \int_{-\infty}^{\infty} \frac{(\min\{1, |x^\delta y|\})^\beta}{|1+x^\delta y|^{\lambda+\beta}} f(x) dx \right]^p dy \\ &\leq K^p(\sigma) \int_{-\infty}^{\infty} |x|^{p(1-\delta\sigma)-1} f^p(x) dx. \end{aligned} \quad (2.10)$$

*Proof.* By Hölder's inequality (cf. [30]) and (2.2), we derive that

$$\begin{aligned}
& \left[ \int_{-\infty}^{\infty} \frac{(\min\{1, |x^\delta y|\})^\beta}{|1 + x^\delta y|^{\lambda+\beta}} f(x) dx \right]^p \\
&= \left\{ \int_{-\infty}^{\infty} \frac{(\min\{1, |x^\delta y|\})^\beta}{|1 + x^\delta y|^{\lambda+\beta}} \left[ \frac{|x|^{(1-\delta\sigma)/q}}{|y|^{(1-\sigma)/p}} f(x) \right] \left[ \frac{|y|^{(1-\sigma)/p}}{|x|^{(1-\delta\sigma)/q}} \right] dx \right\}^p \\
&\leq \int_{-\infty}^{\infty} \frac{(\min\{1, |x^\delta y|\})^\beta}{|1 + x^\delta y|^{\lambda+\beta}} \frac{|x|^{(1-\delta\sigma)(p-1)}}{|y|^{1-\sigma}} f^p(x) dx \\
&\quad \times \left[ \int_{-\infty}^{\infty} \frac{(\min\{1, |x^\delta y|\})^\beta}{|1 + x^\delta y|^{\lambda+\beta}} \frac{|y|^{(1-\sigma)(q-1)}}{|x|^{1-\delta\sigma}} dx \right]^{p-1} \\
&= \frac{(\omega(\sigma, y))^{p-1}}{|y|^{p\sigma-1}} \int_{-\infty}^{\infty} \frac{(\min\{1, |x^\delta y|\})^\beta}{|1 + x^\delta y|^{\lambda+\beta}} \frac{|x|^{(1-\delta\sigma)(p-1)}}{|y|^{1-\sigma}} f^p(x) dx.
\end{aligned} \tag{2.11}$$

Then, by (2.4) and Fubini's theorem (cf. [31]), it follows that

$$\begin{aligned}
J &\leq K^{p-1}(\sigma) \int_{-\infty}^{\infty} \left[ \int_{-\infty}^{\infty} \frac{(\min\{1, |x^\delta y|\})^\beta}{|1 + x^\delta y|^{\lambda+\beta}} \frac{|x|^{(1-\delta\sigma)(p-1)}}{|y|^{1-\sigma}} f^p(x) dx \right] dy \\
&= K^{p-1}(\sigma) \int_{-\infty}^{\infty} \varpi(\sigma, x) |x|^{p(1-\delta\sigma)-1} f^p(x) dx.
\end{aligned} \tag{2.12}$$

Hence, by (2.4), inequality (2.10) follows.  $\square$

### 3 Main Results and Applications

**Theorem 3.1.** Let  $p > 1, \frac{1}{p} + \frac{1}{q} = 1, \beta > -1, \mu, \sigma > -\beta, \mu + \sigma = \lambda < 1 - \beta, \delta \in \{-1, 1\}, K(\sigma)$  as indicated by (2.4) (or (2.9)).

If  $f, g \geq 0$ , satisfy

$$0 < \int_{-\infty}^{\infty} |x|^{p(1-\delta\sigma)-1} f^p(x) dx < \infty$$

and

$$0 < \int_{-\infty}^{\infty} |y|^{q(1-\sigma)-1} g^q(y) dy < \infty,$$

then we have the following equivalent inequalities:

$$\begin{aligned}
I &: = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{(\min\{1, |x^\delta y|\})^\beta}{|1 + x^\delta y|^{\lambda+\beta}} f(x) g(y) dx dy \\
&< K(\sigma) \left[ \int_{-\infty}^{\infty} |x|^{p(1-\delta\sigma)-1} f^p(x) dx \right]^{\frac{1}{p}} \left[ \int_{-\infty}^{\infty} |y|^{q(1-\sigma)-1} g^q(y) dy \right]^{\frac{1}{q}},
\end{aligned} \tag{3.1}$$

$$\begin{aligned}
J &: = \int_{-\infty}^{\infty} |y|^{p\sigma-1} \left[ \int_{-\infty}^{\infty} \frac{(\min\{1, |x^\delta y|\})^\beta}{|1 + x^\delta y|^{\lambda+\beta}} f(x) dx \right]^p dy \\
&< K^p(\sigma) \int_{-\infty}^{\infty} |x|^{p(1-\delta\sigma)-1} f^p(x) dx.
\end{aligned} \tag{3.2}$$

where, the constant factors  $K(\sigma)$  and  $K^p(\sigma)$  are the best possible.

In particular, for  $\delta = 1$ , we obtain the following equivalent inequalities:

$$\begin{aligned} & \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{(\min\{1, |xy|\})^\beta}{|1+xy|^{\lambda+\beta}} f(x)g(y) dx dy \\ & < K(\sigma) \left[ \int_{-\infty}^{\infty} |x|^{p(1-\sigma)-1} f^p(x) dx \right]^{\frac{1}{p}} \left[ \int_{-\infty}^{\infty} |y|^{q(1-\sigma)-1} g^q(y) dy \right]^{\frac{1}{q}}, \end{aligned} \quad (3.3)$$

$$\begin{aligned} & \int_{-\infty}^{\infty} |y|^{p\sigma-1} \left[ \int_{-\infty}^{\infty} \frac{(\min\{1, |xy|\})^\beta}{|1+xy|^{\lambda+\beta}} f(x) dx \right]^p dy \\ & < K^p(\sigma) \int_{-\infty}^{\infty} |x|^{p(1-\sigma)-1} f^p(x) dx. \end{aligned} \quad (3.4)$$

*Proof.* If (2.11) takes the form of equality for some  $y \in (-\infty, 0) \cup (0, \infty)$ , then, there exist constants  $A$  and  $B$ , such that they are not all zero, and

$$A \frac{|x|^{(1-\delta\sigma)(p-1)}}{|y|^{1-\sigma}} f^p(x) = B \frac{|y|^{(1-\sigma)(q-1)}}{|x|^{1-\delta\sigma}} \text{ a.e. in } \mathbf{R}.$$

Let us suppose that  $A \neq 0$  (otherwise  $B = A = 0$ ). Then it follows that

$$|x|^{p(1-\delta\sigma)-1} f^p(x) = |y|^{q(1-\sigma)} \frac{B}{A|x|} \text{ a.e. in } \mathbf{R},$$

which contradicts the fact that

$$0 < \int_{-\infty}^{\infty} |x|^{p(1-\delta\sigma)-1} f^p(x) dx < \infty.$$

Hence (2.11) takes the form of strict inequality. So does (2.12), and we obtain (3.2).

By Hölder's inequality (cf. [30]), we have

$$\begin{aligned} I &= \int_{-\infty}^{\infty} \left[ |y|^{\sigma-\frac{1}{p}} \int_{-\infty}^{\infty} \frac{(\min\{1, |x^\delta y|\})^\beta}{|1+x^\delta y|^{\lambda+\beta}} f(x) dx \right] (|y|^{\frac{1}{p}-\sigma} g(y)) dy \\ &\leq J^{\frac{1}{p}} \left[ \int_{-\infty}^{\infty} |y|^{q(1-\sigma)-1} g^q(y) dy \right]^{\frac{1}{q}}. \end{aligned} \quad (3.5)$$

Then by (3.2), we get (3.1). On the other hand, suppose that (3.1) is valid. We then set

$$g(y) := |y|^{p\sigma-1} \left[ \int_{-\infty}^{\infty} \frac{(\min\{1, |x^\delta y|\})^\beta}{|1+x^\delta y|^{\lambda+\beta}} f(x) dx \right]^{p-1} \quad (y \neq 0), \quad (3.6)$$

and then

$$J = \int_{-\infty}^{\infty} |y|^{q(1-\sigma)-1} g^q(y) dy.$$



By (2.12), we have  $J < \infty$ . If  $J = 0$ , then (3.2) is trivially true; if  $0 < J < \infty$ , then by (3.1), we obtain

$$\begin{aligned} 0 &< \int_{-\infty}^{\infty} |y|^{q(1-\sigma)-1} g^q(y) dy = J = I \\ &< K(\sigma) \left[ \int_{-\infty}^{\infty} |x|^{p(1-\delta\sigma)-1} f^p(x) dx \right]^{\frac{1}{p}} \left[ \int_{-\infty}^{\infty} |y|^{q(1-\sigma)-1} g^q(y) dy \right]^{\frac{1}{q}} < \infty, \end{aligned} \quad (3.7)$$

$$J^{\frac{1}{p}} = \left[ \int_{-\infty}^{\infty} |y|^{q(1-\sigma)-1} g^q(y) dy \right]^{\frac{1}{p}} < K(\sigma) \left[ \int_{-\infty}^{\infty} |x|^{p(1-\delta\sigma)-1} f^p(x) dx \right]^{\frac{1}{p}}. \quad (3.8)$$

Hence, we obtain (3.2), which is equivalent to (3.1).

We set  $E_{\delta} := \{x \in \mathbf{R}; |x|^{\delta} \geq 1\}$ , and  $E_{\delta}^+ := E_{\delta} \cap \mathbf{R}_+ = \{x \in \mathbf{R}_+; x^{\delta} \geq 1\}$ . For  $\varepsilon > 0$ , we define two functions  $\tilde{f}(x), \tilde{g}(y)$  as follows:

$$\begin{aligned} \tilde{f}(x) &: = \begin{cases} |x|^{\delta(\sigma-\frac{2\varepsilon}{p})-1}, & x \in E_{\delta} \\ 0, & x \in \mathbf{R} \setminus E_{\delta} \end{cases}, \\ \tilde{g}(y) &: = \begin{cases} 0, & y \in (-\infty, -1] \cup [1, \infty) \\ |y|^{\sigma+\frac{2\varepsilon}{q}-1}, & y \in (-1, 1) \end{cases}. \end{aligned}$$

Then we obtain

$$\begin{aligned} \tilde{L} &: = \left[ \int_{-\infty}^{\infty} |x|^{p(1-\delta\sigma)-1} \tilde{f}^p(x) dx \right]^{\frac{1}{p}} \left[ \int_{-\infty}^{\infty} |y|^{q(1-\sigma)-1} \tilde{g}^q(y) dy \right]^{\frac{1}{q}} \\ &= 2 \left( \int_{E_{\delta}^+} x^{-2\delta\varepsilon-1} dx \right)^{\frac{1}{p}} \left( \int_0^1 y^{2\varepsilon-1} dy \right)^{\frac{1}{q}} = \frac{1}{\varepsilon}. \end{aligned}$$

We find

$$\begin{aligned} I(x) &: = \int_{-1}^1 \frac{(\max\{1, |x^{\delta}y|\})^{\beta}}{|1+x^{\delta}y|^{\lambda+\beta}} |y|^{\sigma+\frac{2\varepsilon}{q}-1} dy \\ &\stackrel{y=-Y}{=} \int_{-1}^1 \frac{(\max\{1, |-x^{\delta}Y|\})^{\beta}}{|1-x^{\delta}Y|^{\lambda+\beta}} |-Y|^{\sigma+\frac{2\varepsilon}{q}-1} dY = I(-x), \end{aligned}$$

and then  $I(x)$  is an even function. It follows that

$$\begin{aligned} \tilde{I} &:= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{(\min\{1, |x^{\delta}y|\})^{\beta}}{|1+x^{\delta}y|^{\lambda+\beta}} \tilde{f}(x) \tilde{g}(y) dx dy \\ &= \int_{E_{\delta}} |x|^{\delta(\sigma-\frac{2\varepsilon}{p})-1} I(x) dx = 2 \int_{E_{\delta}^+} x^{\delta(\sigma-\frac{2\varepsilon}{p})-1} I(x) dx \\ &\stackrel{u=x^{\delta}y}{=} 2 \int_{E_{\delta}^+} x^{-2\delta\varepsilon-1} \left[ \int_{-x^{\delta}}^{x^{\delta}} \frac{(\min\{1, |u|\})^{\beta}}{|1+u|^{\lambda+\beta}} |u|^{\sigma+\frac{2\varepsilon}{q}-1} du \right] dx. \end{aligned}$$

Setting  $v = x^\delta$  in the above integral, by Fubini's theorem (cf. [31]), we find

$$\begin{aligned}
\tilde{I} &= 2 \int_1^\infty v^{-2\varepsilon-1} \left[ \int_{-v}^v \frac{(\min\{1, |u|\})^\beta}{|1+u|^{\lambda+\beta}} |u|^{\sigma+\frac{2\varepsilon}{q}-1} du \right] dv \\
&= 2 \int_1^\infty v^{-2\varepsilon-1} \left\{ \int_0^v \left[ \frac{(\min\{1, u\})^\beta}{|1-u|^{\lambda+\beta}} + \frac{(\min\{1, u\})^\beta}{(1+u)^{\lambda+\beta}} \right] u^{\sigma+\frac{2\varepsilon}{q}-1} du \right\} dv \\
&= 2 \int_1^\infty v^{-2\varepsilon-1} \left\{ \int_0^1 \left[ \frac{1}{(1-u)^{\lambda+\beta}} + \frac{1}{(1+u)^{\lambda+\beta}} \right] u^{\beta+\sigma+\frac{2\varepsilon}{q}-1} du \right\} dv \\
&\quad + 2 \int_1^\infty v^{-2\varepsilon-1} \left\{ \int_1^v \left[ \frac{1}{(u-1)^{\lambda+\beta}} + \frac{1}{(1+u)^{\lambda+\beta}} \right] u^{\sigma+\frac{2\varepsilon}{q}-1} du \right\} dv \\
&= \frac{1}{\varepsilon} \int_0^1 \left[ \frac{1}{(1-u)^{\lambda+\beta}} + \frac{1}{(1+u)^{\lambda+\beta}} \right] u^{\beta+\sigma+\frac{2\varepsilon}{q}-1} du \\
&\quad + 2 \int_1^\infty \left( \int_u^\infty v^{-2\varepsilon-1} dv \right) \left[ \frac{1}{(u-1)^{\lambda+\beta}} + \frac{1}{(1+u)^{\lambda+\beta}} \right] u^{\sigma+\frac{2\varepsilon}{q}-1} du \\
&= \frac{1}{\varepsilon} \left\{ \int_0^1 \left[ \frac{1}{(1-u)^{\lambda+\beta}} + \frac{1}{(1+u)^{\lambda+\beta}} \right] u^{\beta+\sigma+\frac{2\varepsilon}{q}-1} du \right. \\
&\quad \left. + \int_1^\infty \left[ \frac{1}{(u-1)^{\lambda+\beta}} + \frac{1}{(1+u)^{\lambda+\beta}} \right] u^{\sigma-\frac{2\varepsilon}{p}-1} du \right\}.
\end{aligned}$$

If the constant factor  $K(\sigma)$  in (3.1) is not the best possible, then, there exists a positive number  $k$ , with  $K(\sigma) < k$ , such that (3.1) is valid when replacing  $K(\sigma)$  by  $k$ . Then in particular, we have  $\varepsilon \tilde{I} < \varepsilon k \tilde{L}$ , and

$$\begin{aligned}
&\int_0^1 \left[ \frac{1}{(1-u)^{\lambda+\beta}} + \frac{1}{(1+u)^{\lambda+\beta}} \right] u^{\beta+\sigma+\frac{2\varepsilon}{q}-1} du \\
&+ \int_1^\infty \left[ \frac{1}{(u-1)^{\lambda+\beta}} + \frac{1}{(1+u)^{\lambda+\beta}} \right] u^{\sigma-\frac{2\varepsilon}{p}-1} du = \varepsilon \tilde{I} < \varepsilon k \tilde{L} = k.
\end{aligned} \tag{3.9}$$

By (2.5), (2.6) and Fatou's lemma (cf. [31]), we have

$$\begin{aligned}
K(\sigma) &= \int_0^1 \left[ \frac{1}{(1-u)^{\lambda+\beta}} + \frac{1}{(1+u)^{\lambda+\beta}} \right] u^{\beta+\sigma-1} du \\
&\quad + \int_1^\infty \left[ \frac{1}{(u-1)^{\lambda+\beta}} + \frac{1}{(1+u)^{\lambda+\beta}} \right] u^{\sigma-1} du \\
&= \int_0^1 \lim_{\varepsilon \rightarrow 0^+} \left[ \frac{1}{(1-u)^{\lambda+\beta}} + \frac{1}{(1+u)^{\lambda+\beta}} \right] u^{\beta+\sigma+\frac{2\varepsilon}{q}-1} du \\
&\quad + \int_1^\infty \lim_{\varepsilon \rightarrow 0^+} \left[ \frac{1}{(u-1)^{\lambda+\beta}} + \frac{1}{(1+u)^{\lambda+\beta}} \right] u^{\sigma-\frac{2\varepsilon}{p}-1} du \\
&\leq \underline{\lim}_{\varepsilon \rightarrow 0^+} \left\{ \int_0^1 \left[ \frac{1}{(1-u)^{\lambda+\beta}} + \frac{1}{(1+u)^{\lambda+\beta}} \right] u^{\beta+\sigma+\frac{2\varepsilon}{q}-1} du \right. \\
&\quad \left. + \int_1^\infty \left[ \frac{1}{(u-1)^{\lambda+\beta}} + \frac{1}{(1+u)^{\lambda+\beta}} \right] u^{\sigma-\frac{2\varepsilon}{p}-1} du \right\} \leq k,
\end{aligned}$$

which contradicts the fact that  $k < K(\sigma)$ . Hence the constant factor  $K(\sigma)$  in (3.1) is the best possible.

If the constant factor in (3.2) is not the best possible, then by (3.5) we would reach the contradiction that the constant factor in (3.1) is not the best possible.  $\square$

**Theorem 3.2.** *If in the assumptions of Theorem 3.1, we replace  $p > 1$  by  $0 < p < 1$ , we obtain the equivalent reverses of (3.1) and (3.2) with the same best constant factors.*

*Proof.* By Hölder's reverse inequality (cf. [30]), we derive the reverses of (2.11), (2.12), (2.10) and (3.5). It is easy to obtain the reverse of (3.2). In view of the reverses of (3.2) and (3.5), we obtain the reverse of (3.1). On the other hand, if we suppose that the reverse of (3.1) is valid, then if we set  $g(y)$  as in (3.6), by the reverse of (2.12), we have  $J > 0$ . If  $J = \infty$ , then the reverse of (3.2) is obviously true; if  $J < \infty$ , then by the reverse of (3.1), we obtain the reverses of (3.7) and (3.8). Hence, we obtain the reverse of (3.2), which is equivalent to the reverse of (3.1).

If the constant factor  $K(\sigma)$  in the reverse of (3.1) is not the best possible, then, there exists a positive constant  $k$ , with  $k > K(\sigma)$ , such that the reverse of (3.1) is still valid when replacing  $K(\sigma)$  by  $k$ . By the reverse of (3.9), we have

$$\begin{aligned} & \int_0^1 \left[ \frac{1}{(1-u)^{\lambda+\beta}} + \frac{1}{(1+u)^{\lambda+\beta}} \right] u^{\beta+\sigma+\frac{2\varepsilon}{q}-1} du \\ & + \int_1^\infty \left[ \frac{1}{(u-1)^{\lambda+\beta}} + \frac{1}{(1+u)^{\lambda+\beta}} \right] u^{\sigma-\frac{2\varepsilon}{p}-1} du > k. \end{aligned} \quad (3.10)$$

By Levi's theorem (cf. [31]), we find

$$\begin{aligned} & \int_1^\infty \left[ \frac{1}{(u-1)^{\lambda+\beta}} + \frac{1}{(1+u)^{\lambda+\beta}} \right] u^{\sigma-\frac{2\varepsilon}{p}-1} du \\ \rightarrow & \int_1^\infty \left[ \frac{1}{(u-1)^{\lambda+\beta}} + \frac{1}{(1+u)^{\lambda+\beta}} \right] u^{\sigma-1} du (\varepsilon \rightarrow 0^+). \end{aligned}$$

There exists a constant  $\delta_0 > 0$ , such that  $\sigma - \frac{1}{2}\delta_0 > -\beta$ , and then  $K(\sigma - \frac{\delta_0}{2}) \in \mathbf{R}_+$ . For  $0 < \varepsilon < \frac{\delta_0|q|}{4}$  ( $q < 0$ ), since  $u^{\beta+\sigma+\frac{2\varepsilon}{q}-1} \leq u^{\beta+\sigma-\frac{\delta_0}{2}-1}$ ,  $u \in (0, 1]$ , and

$$0 < \int_0^1 \left[ \frac{1}{(1-u)^{\lambda+\beta}} + \frac{1}{(1+u)^{\lambda+\beta}} \right] u^{\beta+\sigma-\frac{\delta_0}{2}-1} du \leq K(\sigma - \frac{\delta_0}{2}),$$

then by Lebesgue's control convergence theorem (cf. [31]), we have

$$\begin{aligned} & \int_0^1 \left[ \frac{1}{(1-u)^{\lambda+\beta}} + \frac{1}{(1+u)^{\lambda+\beta}} \right] u^{\beta+\sigma+\frac{2\varepsilon}{q}-1} du \\ \rightarrow & \int_0^1 \left[ \frac{1}{(1-u)^{\lambda+\beta}} + \frac{1}{(1+u)^{\lambda+\beta}} \right] u^{\beta+\sigma-1} du (\varepsilon \rightarrow 0^+). \end{aligned}$$

By (3.10) and the above results, for  $\varepsilon \rightarrow 0^+$ , we get  $K(\sigma) \geq k$ , which contradicts the fact that  $k > K(\sigma)$ . Hence, the constant factor  $K(\sigma)$  in the reverse of (3.1) is the best possible.

If the constant factor in the reverse of (3.2) is not the best possible, then, by the reverse of (3.5), we would reach the contradiction that the constant factor in the reverse of (3.1) is not the best possible.  $\square$

*Remark 3.3.* (i) For  $\delta = -1$  in (3.1) and (3.2), replacing  $|x|^\lambda f(x)$  by  $f(x)$ , we obtain  $0 < \int_{-\infty}^{\infty} |x|^{p(1-\mu)-1} f^p(x) dx < \infty$ , and the following equivalent inequalities with the homogeneous kernel and the best possible constant factors:

$$\begin{aligned} & \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{(\min\{|x|, |y|\})^\beta}{|x+y|^{\lambda+\beta}} f(x)g(y) dx dy \\ & < K(\sigma) \left[ \int_{-\infty}^{\infty} |x|^{p(1-\mu)-1} f^p(x) dx \right]^{\frac{1}{p}} \left[ \int_{-\infty}^{\infty} |y|^{q(1-\sigma)-1} g^q(y) dy \right]^{\frac{1}{q}}, \end{aligned} \quad (3.11)$$

$$\begin{aligned} & \int_{-\infty}^{\infty} |y|^{p\sigma-1} \left[ \int_{-\infty}^{\infty} \frac{(\max\{|x|, |y|\})^\beta}{|x+y|^{\lambda+\beta}} f(x) dx \right]^p dy \\ & < K^p(\sigma) \int_{-\infty}^{\infty} |x|^{p(1-\mu)-1} f^p(x) dx. \end{aligned} \quad (3.12)$$

In particular, for  $\lambda = 0 = \mu + \sigma$  ( $\mu, \sigma > -\beta$ ),  $0 < \beta < 1$ , we find

$$\begin{aligned} K(\sigma) &= K_0(\sigma) := \frac{1}{\beta + \sigma} F(\beta, \beta + \sigma, 1 + \beta + \sigma, -1) \\ &\quad + \frac{1}{\beta + \mu} F(\beta, \beta + \mu, 1 + \beta + \mu, -1) \\ &\quad + B(1 - \beta, \beta + \sigma) + B(1 - \beta, \beta + \mu), \end{aligned} \quad (3.13)$$

and the following equivalent inequalities:

$$\begin{aligned} & \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \left( \frac{\min\{|x|, |y|\}}{|x+y|} \right)^\beta f(x)g(y) dx dy \\ & < K_0(\sigma) \left[ \int_{-\infty}^{\infty} |x|^{p(1-\mu)-1} f^p(x) dx \right]^{\frac{1}{p}} \left[ \int_{-\infty}^{\infty} |y|^{q(1-\sigma)-1} g^q(y) dy \right]^{\frac{1}{q}}, \end{aligned} \quad (3.14)$$

$$\begin{aligned} & \int_{-\infty}^{\infty} |y|^{p\sigma-1} \left[ \int_{-\infty}^{\infty} \left( \frac{\min\{|x|, |y|\}}{|x+y|} \right)^\beta f(x) dx \right]^p dy \\ & < K_0^p(\sigma) \int_{-\infty}^{\infty} |x|^{p(1-\mu)-1} f^p(x) dx. \end{aligned} \quad (3.15)$$

(ii) For  $\lambda = 0 = \mu + \sigma$  ( $\mu, \sigma > -\beta$ ),  $0 < \beta < 1$  in (3.1) and (3.2), we have the following equivalent inequalities:

$$\begin{aligned} & \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \left( \frac{\min\{1, |x^\delta y|\}}{|1 + x^\delta y|} \right)^\beta f(x)g(y) dx dy \\ & < K_0(\sigma) \left[ \int_{-\infty}^{\infty} |x|^{p(1-\delta\sigma)-1} f^p(x) dx \right]^{\frac{1}{p}} \left[ \int_{-\infty}^{\infty} |y|^{q(1-\sigma)-1} g^q(y) dy \right]^{\frac{1}{q}}, \end{aligned} \quad (3.16)$$

$$\begin{aligned}
& \int_{-\infty}^{\infty} |y|^{p\sigma-1} \left[ \int_{-\infty}^{\infty} \left( \frac{\min\{1, |x^\delta y|\}}{|1+x^\delta y|} \right)^\beta f(x) dx \right]^p dy \\
& < K_0^p(\sigma) \int_{-\infty}^{\infty} |x|^{p(1-\delta\sigma)-1} f^p(x) dx.
\end{aligned} \tag{3.17}$$

In particular, for  $\delta = 1$ , we have the following equivalent inequalities (cf. [25], for  $\sigma = \mu = 0$ ):

$$\begin{aligned}
& \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \left( \frac{\min\{1, |xy|\}}{|1+xy|} \right)^\beta f(x) g(y) dx dy \\
& < K_0(\sigma) \left[ \int_{-\infty}^{\infty} |x|^{p(1-\sigma)-1} f^p(x) dx \right]^{\frac{1}{p}} \left[ \int_{-\infty}^{\infty} |y|^{q(1-\sigma)-1} g^q(y) dy \right]^{\frac{1}{q}},
\end{aligned} \tag{3.18}$$

$$\begin{aligned}
& \int_{-\infty}^{\infty} |y|^{p\sigma-1} \left[ \int_{-\infty}^{\infty} \left( \frac{\min\{1, |xy|\}}{|1+xy|} \right)^\beta f(x) dx \right]^p dy \\
& < K_0^p(\sigma) \int_{-\infty}^{\infty} |x|^{p(1-\sigma)-1} f^p(x) dx.
\end{aligned} \tag{3.19}$$

(iii) For  $\beta = 0 < \lambda < 1$ ,  $\sigma = \mu = \frac{\lambda}{2}$  in (3.3), we obtain

$$K\left(\frac{\lambda}{2}\right) = \int_0^\infty \frac{u^{\frac{\lambda}{2}-1}}{(1+u)^\lambda} du + 2 \int_0^1 \frac{u^{\frac{\lambda}{2}-1}}{(1-u)^\lambda} du = k_\lambda,$$

and then (1.3) follows. Hence, (3.1)-(3.3) is an extension of (1.3).

## 4 Some Corollaries

In the following two sections, if the constant factors are related to  $K_1(\sigma)$ , then we call them Hardy-type inequalities (operators) of the first kind; if the constant factors are related to  $K_2(\sigma)$ , then we call them Hardy-type inequalities (operators) of the second kind.

Setting the kernel

$$H(xy) := \begin{cases} 0, & |xy| > 1 \\ \frac{(\min\{1, |xy|\})^\beta}{|1+xy|^{\lambda+\beta}}, & |xy| \leq 1 \end{cases},$$

it follows that

$$\begin{aligned}
H(u) &= \begin{cases} 0, & |u| > 1 \\ \frac{(\min\{1, |u|\})^\beta}{|1+u|^{\lambda+\beta}}, & |u| \leq 1 \end{cases}, \\
\int_{-\infty}^{\infty} H(u) |u|^{\sigma-1} du &= \int_{-1}^1 \frac{(\min\{1, |u|\})^\beta}{|1+u|^{\lambda+\beta}} |u|^{\sigma-1} du = K_1(\sigma).
\end{aligned}$$

In view of Theorems 3.1-3.2 (for  $\delta = 1$ ), we obtain the following Hardy-type inequalities of the first kind with the non-homogeneous kernel:

**Corollary 4.1.** Suppose that  $p > 1, \frac{1}{p} + \frac{1}{q} = 1, \beta > -1, \mu, \sigma > -\beta, \mu + \sigma = \lambda < 1 - \beta, K_1(\sigma)$  is indicated by (2.5) (or 2.7). If  $f, g \geq 0$ , satisfy

$$0 < \int_{-\infty}^{\infty} |x|^{p(1-\sigma)-1} f^p(x) dx < \infty$$

and

$$0 < \int_{-\infty}^{\infty} |y|^{q(1-\sigma)-1} g^q(y) dy < \infty,$$

then we have the following equivalent inequalities:

$$\begin{aligned} & \int_{-\infty}^{\infty} \left[ \int_{-\frac{1}{|y|}}^{\frac{1}{|y|}} \frac{(\min\{1, |xy|\})^\beta}{|1+xy|^{\lambda+\beta}} f(x) dx \right] g(y) dy \\ & < K_1(\sigma) \left[ \int_{-\infty}^{\infty} |x|^{p(1-\sigma)-1} f^p(x) dx \right]^{\frac{1}{p}} \left[ \int_{-\infty}^{\infty} |y|^{q(1-\sigma)-1} g^q(y) dy \right]^{\frac{1}{q}}, \end{aligned} \quad (4.1)$$

$$\begin{aligned} & \int_{-\infty}^{\infty} |y|^{p\sigma-1} \left[ \int_{-\frac{1}{|y|}}^{\frac{1}{|y|}} \frac{(\min\{1, |xy|\})^\beta}{|1+x^\delta y|^{\lambda+\beta}} f(x) dx \right]^p dy \\ & < K_1^p(\sigma) \int_{-\infty}^{\infty} |x|^{p(1-\sigma)-1} f^p(x) dx. \end{aligned} \quad (4.2)$$

where, the constant factors  $K_1(\sigma)$  and  $K_1^p(\sigma)$  are the best possible. Replacing  $p > 1$  by  $0 < p < 1$ , we have the equivalent reverses of (4.1) and (4.2) with the same best constant factors.

If we set  $E_y := \{x \in \mathbf{R}; |xy| \geq 1\}$ , and

$$H(xy) := \begin{cases} 0, & |xy| < 1 \\ \frac{(\min\{1, |xy|\})^\beta}{|1+xy|^{\lambda+\beta}}, & |xy| \geq 1 \end{cases},$$

then it follows that

$$\begin{aligned} H(u) &= \begin{cases} 0, & |u| < 1 \\ \frac{(\min\{1, |u|\})^\beta}{|1+u|^{\lambda+\beta}}, & |u| \geq 1 \end{cases}, \\ \int_{-\infty}^{\infty} H(u) |u|^{\sigma-1} du &= \int_{E_1} \frac{(\min\{1, |u|\})^\beta}{|1+u|^{\lambda+\beta}} |u|^{\sigma-1} du = K_2(\sigma). \end{aligned}$$

In view of Theorems 3.1-3.2 (for  $\delta = 1$ ), we have the following Hardy-type inequalities of the second kind with the non-homogeneous kernel:

**Corollary 4.2.** Suppose that  $p > 1, \frac{1}{p} + \frac{1}{q} = 1, \beta > -1, \mu, \sigma > -\beta, \mu + \sigma = \lambda < 1 - \beta, K_2(\sigma)$  is indicated by (2.6) (or 2.8).

If  $f, g \geq 0$ , satisfy

$$0 < \int_{-\infty}^{\infty} |x|^{p(1-\sigma)-1} f^p(x) dx < \infty$$

and

$$0 < \int_{-\infty}^{\infty} |y|^{q(1-\sigma)-1} g^q(y) dy < \infty,$$

then we have the following equivalent inequalities:

$$\begin{aligned} & \int_{-\infty}^{\infty} \left[ \int_{E_y} \frac{(\min\{1, |xy|\})^\beta}{|1+xy|^{\lambda+\beta}} f(x) dx \right] g(y) dy \\ & < K_2(\sigma) \left[ \int_{-\infty}^{\infty} |x|^{p(1-\sigma)-1} f^p(x) dx \right]^{\frac{1}{p}} \left[ \int_{-\infty}^{\infty} |y|^{q(1-\sigma)-1} g^q(y) dy \right]^{\frac{1}{q}}, \end{aligned} \quad (4.3)$$

$$\begin{aligned} & \int_{-\infty}^{\infty} |y|^{p\sigma-1} \left[ \int_{E_y} \frac{(\min\{1, |xy|\})^\beta}{|1+xy|^{\lambda+\beta}} f(x) dx \right]^p dy \\ & < K_2^p(\sigma) \int_{-\infty}^{\infty} |x|^{p(1-\delta\sigma)-1} f^p(x) dx. \end{aligned} \quad (4.4)$$

where, the constant factors  $K_2(\sigma)$  and  $K_2^p(\sigma)$  are the best possible. Replacing  $p > 1$  by  $0 < p < 1$ , we obtain the equivalent reverses of (4.3) and (4.4) with the same best constant factors.

If we set  $\tilde{E}_y := \{x \in \mathbf{R}; |\frac{y}{x}| \leq 1\}$  and

$$K_\lambda(x, y) := \begin{cases} 0, & |\frac{y}{x}| > 1 \\ \frac{(\min\{|x|, |y|\})^\beta}{|x+y|^{\lambda+\beta}}, & |\frac{y}{x}| \leq 1 \end{cases},$$

then it follows

$$\begin{aligned} K_\lambda(1, u) &= \begin{cases} 0, & |u| > 1 \\ \frac{(\min\{1, |u|\})^\beta}{|1+u|^{\lambda+\beta}}, & |u| \leq 1 \end{cases}, \\ \int_{-\infty}^{\infty} K_\lambda(1, u) |u|^{\sigma-1} du &= \int_{-1}^1 \frac{(\min\{1, |u|\})^\beta}{|1+u|^{\lambda+\beta}} |u|^{\sigma-1} du = K_1(\sigma). \end{aligned}$$

In view of Remark 3.3 (i), we have the following Hardy-type inequalities of the first kind with the homogeneous kernel:

**Corollary 4.3.** Suppose that  $p > 1, \frac{1}{p} + \frac{1}{q} = 1, \beta > -1, \mu, \sigma > -\beta, \mu + \sigma = \lambda < 1 - \beta, K_1(\sigma)$  is indicated by (2.5) (or 2.7). If  $f, g \geq 0$ , satisfy

$$0 < \int_{-\infty}^{\infty} |x|^{p(1-\mu)-1} f^p(x) dx < \infty$$

and

$$0 < \int_{-\infty}^{\infty} |y|^{q(1-\sigma)-1} g^q(y) dy < \infty,$$

then we have the following equivalent inequalities:

$$\begin{aligned} & \int_{-\infty}^{\infty} \left[ \int_{\tilde{E}_y} \frac{(\min\{|x|, |y|\})^\beta}{|x+y|^{\lambda+\beta}} f(x) dx \right] g(y) dy \\ & < K_1(\sigma) \left[ \int_{-\infty}^{\infty} |x|^{p(1-\mu)-1} f^p(x) dx \right]^{\frac{1}{p}} \left[ \int_{-\infty}^{\infty} |y|^{q(1-\sigma)-1} g^q(y) dy \right]^{\frac{1}{q}}, \end{aligned} \quad (4.5)$$

$$\begin{aligned}
& \int_{-\infty}^{\infty} |y|^{p\sigma-1} \left[ \int_{\tilde{E}_y} \frac{(\min\{|x|, |y|\})^\beta}{|x+y|^{\lambda+\beta}} f(x) dx \right]^p dy \\
& < K_1^p(\sigma) \int_{-\infty}^{\infty} |x|^{p(1-\mu)-1} f^p(x) dx.
\end{aligned} \tag{4.6}$$

where the constant factors  $K_1(\sigma)$  and  $K_1^p(\sigma)$  are the best possible. Replacing  $p > 1$  by  $0 < p < 1$ , we derive the equivalent reverses of (4.5) and (4.6) with the same best constant factors.

Setting the kernel

$$K_\lambda(x, y) := \begin{cases} 0, & |\frac{y}{x}| < 1 \\ \frac{(\min\{|x|, |y|\})^\beta}{|x+y|^{\lambda+\beta}}, & |\frac{y}{x}| \geq 1 \end{cases},$$

then it follows that

$$\begin{aligned}
K_\lambda(1, u) &= \begin{cases} 0, & |u| < 1 \\ \frac{(\min\{1, |u|\})^\beta}{|1+u|^{\lambda+\beta}}, & |u| \geq 1 \end{cases}, \\
\int_{-\infty}^{\infty} K_\lambda(1, u) |u|^{\sigma-1} du &= \int_{E_1} \frac{(\min\{1, |u|\})^\beta}{|1+u|^{\lambda+\beta}} |u|^{\sigma-1} du = K_2(\sigma).
\end{aligned}$$

In view of Remark 3.3 (i), we have the following Hardy-type inequalities of the second kind with the homogeneous kernel:

**Corollary 4.4.** Suppose that  $p > 1, \frac{1}{p} + \frac{1}{q} = 1, \beta < 1, \mu, \sigma > 0, \mu + \sigma = \lambda < 1 - \beta, K_2(\sigma)$  is indicated by (2.6) (or 2.8). If  $f, g \geq 0$ , satisfy

$$0 < \int_{-\infty}^{\infty} |x|^{p(1-\mu)-1} f^p(x) dx < \infty$$

and

$$0 < \int_{-\infty}^{\infty} |y|^{q(1-\sigma)-1} g^q(y) dy < \infty,$$

then we have the following equivalent inequalities:

$$\begin{aligned}
& \int_{-\infty}^{\infty} \left[ \int_{-|y|}^{|y|} \frac{(\min\{|x|, |y|\})^\beta}{|x+y|^{\lambda+\beta}} f(x) dx \right] g(y) dy \\
& < K_2(\sigma) \left[ \int_{-\infty}^{\infty} |x|^{p(1-\mu)-1} f^p(x) dx \right]^{\frac{1}{p}} \left[ \int_{-\infty}^{\infty} |y|^{q(1-\sigma)-1} g^q(y) dy \right]^{\frac{1}{q}},
\end{aligned} \tag{4.7}$$

$$\begin{aligned}
& \int_{-\infty}^{\infty} |y|^{p\sigma-1} \left[ \int_{-|y|}^{|y|} \frac{(\min\{|x|, |y|\})^\beta}{|x+y|^{\lambda+\beta}} f(x) dx \right]^p dy \\
& < K_2^p(\sigma) \int_{-\infty}^{\infty} |x|^{p(1-\mu)-1} f^p(x) dx.
\end{aligned} \tag{4.8}$$

where the constant factors  $K_2(\sigma)$  and  $K_2^p(\sigma)$  are the best possible. Replacing  $p > 1$  by  $0 < p < 1$ , we get the equivalent reverses of (4.7) and (4.8) with the same best constant factors.



## 5 Operator Expressions

Suppose that  $p > 1, \frac{1}{p} + \frac{1}{q} = 1, \beta > -1, \mu, \sigma > -\beta, \mu + \sigma = \lambda < 1 - \beta$ . We set the following functions:  $\varphi(x) := |x|^{p(1-\sigma)-1}, \psi(y) := |y|^{q(1-\sigma)-1}, \phi(x) := |x|^{p(1-\mu)-1} (x, y \in \mathbf{R})$ , wherefrom,  $\psi^{1-p}(y) = |y|^{p\sigma-1}$ . Define the following real normed linear space:

$$L_{p,\varphi}(\mathbf{R}) := \left\{ f : \|f\|_{p,\varphi} := \left( \int_{-\infty}^{\infty} \varphi(x) |f(x)|^p dx \right)^{\frac{1}{p}} < \infty \right\},$$

wherefrom,

$$\begin{aligned} L_{p,\psi^{1-p}}(\mathbf{R}) &= \left\{ h : \|h\|_{p,\psi^{1-p}} := \left( \int_{-\infty}^{\infty} \psi^{1-p}(y) |h(y)|^p dy \right)^{\frac{1}{p}} < \infty \right\}, \\ L_{p,\phi}(\mathbf{R}) &= \left\{ g : \|g\|_{p,\phi} := \left( \int_{-\infty}^{\infty} \phi(x) |g(x)|^p dx \right)^{\frac{1}{p}} < \infty \right\}. \end{aligned}$$

(a) In view of Theorem 3.1 ( $\delta = 1$ ), for  $f \in L_{p,\varphi}(\mathbf{R})$ , setting

$$H^{(1)}(y) := \int_{-\infty}^{\infty} \frac{(\min\{1, |xy|\})^\beta}{|1+xy|^{\lambda+\beta}} |f(x)| dx \quad (y \in \mathbf{R}),$$

by (3.4), we have

$$\|H^{(1)}\|_{p,\psi^{1-p}} = \left[ \int_{-\infty}^{\infty} \psi^{1-p}(y) (H^{(1)}(y))^p dy \right]^{\frac{1}{p}} < K(\sigma) \|f\|_{p,\varphi} < \infty. \quad (5.1)$$

**Definition 5.1.** Define the Hilbert-type integral operator with the non-homogeneous kernel in the whole plane  $T^{(1)} : L_{p,\varphi}(\mathbf{R}) \rightarrow L_{p,\psi^{1-p}}(\mathbf{R})$  as follows: For any  $f \in L_{p,\varphi}(\mathbf{R})$ , there exists a unique representation  $T^{(1)}f = H^{(1)} \in L_{p,\psi^{1-p}}(\mathbf{R})$ , satisfying

$$T^{(1)}f(y) = H^{(1)}(y),$$

for any  $y \in \mathbf{R}$ .

In view of (5.1), it follows that  $\|T^{(1)}f\|_{p,\psi^{1-p}} = \|H^{(1)}\|_{p,\psi^{1-p}} \leq K(\sigma) \|f\|_{p,\varphi}$ . Then, the operator  $T^{(1)}$  is bounded satisfying

$$\|T^{(1)}\| = \sup_{f(\neq 0) \in L_{p,\varphi}(\mathbf{R}_+)} \frac{\|T^{(1)}f\|_{p,\psi^{1-p}}}{\|f\|_{p,\varphi}} \leq K(\sigma).$$

Since the constant factor  $K(\sigma)$  in (5.1) is the best possible, we have  $\|T^{(1)}\| = K(\sigma)$ .

If we define the formal inner product of  $T^{(1)}f$  and  $g$  as follows:

$$\begin{aligned} (T^{(1)}f, g) &: = \int_{-\infty}^{\infty} \left[ \int_{-\infty}^{\infty} \frac{(\min\{1, |xy|\})^\beta}{|1+xy|^{\lambda+\beta}} f(x) dx \right] g(y) dy \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{(\min\{1, |xy|\})^\beta}{|1+xy|^{\lambda+\beta}} f(x) g(y) dx dy, \end{aligned}$$

then we can rewrite (3.3) and (3.4) in the form:

$$(T^{(1)}f, g) < \|T^{(1)}\| \cdot \|f\|_{p, \Phi} \|g\|_{q, \Psi}, \|T^{(1)}f\|_{p, \Psi^{1-p}} < \|T^{(1)}\| \cdot \|f\|_{p, \Phi}. \quad (5.2)$$

(b) In view of Corollary 4.1, for  $f \in L_{p, \Phi}(\mathbf{R})$ , setting

$$H_1^{(1)}(y) := \int_{-\frac{1}{|y|}}^{\frac{1}{|y|}} \frac{(\min\{1, |xy|\})^\beta}{|1 + xy|^{\lambda+\beta}} |f(x)| dx \quad (y \in \mathbf{R}),$$

by (3.14), we obtain

$$\|H_1^{(1)}\|_{p, \Psi^{1-p}} = \left[ \int_{-\frac{1}{|y|}}^{\frac{1}{|y|}} \Psi^{1-p}(y) (H_1^{(1)}(y))^p dy \right]^{\frac{1}{p}} < K_1(\sigma) \|f\|_{p, \Phi} < \infty. \quad (5.3)$$

**Definition 5.2.** Define the Hilbert-type integral operator of the first kind with the non-homogeneous kernel in the whole plane  $T_1^{(1)} : L_{p, \Phi}(\mathbf{R}) \rightarrow L_{p, \Psi^{1-p}}(\mathbf{R})$  as follows: For any  $f \in L_{p, \Phi}(\mathbf{R})$ , there exists a unique representation  $T_1^{(1)}f = H_1^{(1)} \in L_{p, \Psi^{1-p}}(\mathbf{R})$ , satisfying

$$T_1^{(1)}f(y) = H_1^{(1)}(y),$$

for any  $y \in \mathbf{R}$ .

In view of (5.3), it follows that  $\|T_1^{(1)}f\|_{p, \Psi^{1-p}} = \|H_1^{(1)}\|_{p, \Psi^{1-p}} \leq K_1(\sigma) \|f\|_{p, \Phi}$ , and then the operator  $T_1^{(1)}$  is bounded satisfying

$$\|T_1^{(1)}\| = \sup_{f(\neq 0) \in L_{p, \Phi}(\mathbf{R}_+)} \frac{\|T_1^{(1)}f\|_{p, \Psi^{1-p}}}{\|f\|_{p, \Phi}} \leq K_1(\sigma).$$

Since the constant factor  $K_1(\sigma)$  in (5.3) is the best possible, we have  $\|T_1^{(1)}\| = K_1(\sigma)$ .

If we define the formal inner product of  $T_1^{(1)}f$  and  $g$  as follows:

$$(T_1^{(1)}f, g) := \int_{-\infty}^{\infty} \left[ \int_{-\frac{1}{|y|}}^{\frac{1}{|y|}} \frac{(\min\{1, |xy|\})^\beta}{|1 + xy|^{\lambda+\beta}} f(x) dx \right] g(y) dy,$$

then we can rewrite (3.13) and (3.14) in the following way:

$$(T_1^{(1)}f, g) < \|T_1^{(1)}\| \cdot \|f\|_{p, \Phi} \|g\|_{q, \Psi}, \|T_1^{(1)}f\|_{p, \Psi^{1-p}} < \|T_1^{(1)}\| \cdot \|f\|_{p, \Phi}. \quad (5.4)$$

(c) In view of Corollary 4.2, for  $f \in L_{p, \Phi}(\mathbf{R})$ , setting

$$H_2^{(1)}(y) := \int_{E_y} \frac{(\min\{1, |xy|\})^\beta}{|1 + xy|^{\lambda+\beta}} |f(x)| dx \quad (y \in \mathbf{R}),$$

by (3.16), we have

$$\|H_2^{(1)}\|_{p, \Psi^{1-p}} = \left[ \int_{E_y} \Psi^{1-p}(y) (H_2^{(1)}(y))^p dy \right]^{\frac{1}{p}} < K_2(\sigma) \|f\|_{p, \Phi} < \infty. \quad (5.5)$$

**Definition 5.3.** Define the Hilbert-type integral operator of the second kind with the non-homogeneous kernel in the whole plane  $T_2^{(1)} : L_{p,\phi}(\mathbf{R}) \rightarrow L_{p,\psi^{1-p}}(\mathbf{R})$  as follows: For any  $f \in L_{p,\phi}(\mathbf{R})$ , there exists a unique representation  $T_2^{(1)} f = H_2^{(1)} \in L_{p,\psi^{1-p}}(\mathbf{R})$ , satisfying

$$T_2^{(1)} f(y) = H_2^{(1)}(y),$$

for any  $y \in \mathbf{R}$ .

In view of (5.5), it follows that  $\|T_2^{(1)} f\|_{p,\psi^{1-p}} = \|H_2^{(1)}\|_{p,\psi^{1-p}} \leq K_2(\sigma) \|f\|_{p,\phi}$ , and then the operator  $T_2^{(1)}$  is bounded satisfying

$$\|T_2^{(1)}\| = \sup_{f(\neq 0) \in L_{p,\phi}(\mathbf{R}_+)} \frac{\|T_2^{(1)} f\|_{p,\psi^{1-p}}}{\|f\|_{p,\phi}} \leq K_2(\sigma).$$

Since the constant factor  $K_2(\sigma)$  in (5.5) is the best possible, we have  $\|T_2^{(1)}\| = K_2(\sigma)$ .

If we define the formal inner product of  $T_2^{(1)} f$  and  $g$  as follows:

$$(T_2^{(1)} f, g) := \int_{-\infty}^{\infty} \left[ \int_{E_y} \frac{(\min\{1, |xy|\})^\beta}{|1+xy|^{\lambda+\beta}} f(x) dx \right] g(y) dy,$$

then we can rewrite (3.15) and (3.16) as shown below:

$$(T_2^{(1)} f, g) < \|T_2^{(1)}\| \cdot \|f\|_{p,\phi} \|g\|_{q,\psi}, \|T_2^{(1)} f\|_{p,\psi^{1-p}} < \|T_2^{(1)}\| \cdot \|f\|_{p,\phi}. \quad (5.6)$$

(d) In view of Remark 3.3 (i), for  $f \in L_{p,\phi}(\mathbf{R})$ , setting

$$H^{(2)}(y) := \int_{-\infty}^{\infty} \frac{(\min\{|x|, |y|\})^\beta}{|x+y|^{\lambda+\beta}} |f(x)| dx \quad (y \in \mathbf{R}),$$

by (3.12), we have

$$\|H^{(2)}\|_{p,\psi^{1-p}} = \left[ \int_{-\infty}^{\infty} \psi^{1-p}(y) (H^{(2)}(y))^p dy \right]^{\frac{1}{p}} < K(\sigma) \|f\|_{p,\phi} < \infty. \quad (5.7)$$

**Definition 5.4.** Define the Hilbert-type integral operator with the homogeneous kernel in the whole plane  $T^{(2)} : L_{p,\phi}(\mathbf{R}) \rightarrow L_{p,\psi^{1-p}}(\mathbf{R})$  as follows: For any  $f \in L_{p,\phi}(\mathbf{R})$ , there exists a unique representation  $T^{(2)} f = H^{(2)} \in L_{p,\psi^{1-p}}(\mathbf{R})$ , satisfying

$$T^{(2)} f(y) = H^{(2)}(y),$$

for any  $y \in \mathbf{R}$ .

In view of (5.7), it follows that  $\|T^{(2)} f\|_{p,\psi^{1-p}} = \|H^{(2)}\|_{p,\psi^{1-p}} \leq K(\sigma) \|f\|_{p,\phi}$ , and then the operator  $T^{(2)}$  is bounded satisfying

$$\|T^{(2)}\| = \sup_{f(\neq 0) \in L_{p,\phi}(\mathbf{R}_+)} \frac{\|T^{(2)} f\|_{p,\psi^{1-p}}}{\|f\|_{p,\phi}} \leq K(\sigma).$$

Since the constant factor  $K(\sigma)$  in (5.7) is the best possible, we have  $\|T^{(2)}\| = K(\sigma)$ .

If we define the formal inner product of  $T^{(2)}f$  and  $g$  as follows:

$$\begin{aligned} (T^{(2)}f, g) &: = \int_{-\infty}^{\infty} \left[ \int_{-\infty}^{\infty} \frac{(\min\{|x|, |y|\})^\beta}{|x+y|^{\lambda+\beta}} f(x) dx \right] g(y) dy \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{(\min\{|x|, |y|\})^\beta}{|x+y|^{\lambda+\beta}} f(x) g(y) dx dy, \end{aligned}$$

then we can rewrite (3.11) and (3.12) as follows:

$$(T^{(2)}f, g) < \|T^{(2)}\| \cdot \|f\|_{p,\phi} \|g\|_{q,\psi}, \|T^{(2)}f\|_{p,\psi^{1-p}} < \|T^{(2)}\| \cdot \|f\|_{p,\phi}. \quad (5.8)$$

(e) In view of Corollary 4.3, for  $f \in L_{p,\phi}(\mathbf{R})$ , setting

$$H_1^{(2)}(y) := \int_{\tilde{E}_y} \frac{(\min\{|x|, |y|\})^\beta}{|x+y|^{\lambda+\beta}} |f(x)| dx \quad (y \in \mathbf{R}),$$

by (3.18), we have

$$\|H_1^{(2)}\|_{p,\psi^{1-p}} = \left[ \int_{\tilde{E}_y} \psi^{1-p}(y) (H_1^{(2)}(y))^p dy \right]^{\frac{1}{p}} < K_1(\sigma) \|f\|_{p,\phi} < \infty. \quad (5.9)$$

**Definition 5.5.** Define the Hilbert-type integral operator of the first kind with the homogeneous kernel in the whole plane  $T_1^{(2)} : L_{p,\phi}(\mathbf{R}) \rightarrow L_{p,\psi^{1-p}}(\mathbf{R})$  as follows: For any  $f \in L_{p,\phi}(\mathbf{R})$ , there exists a unique representation  $T_1^{(2)}f = H_1^{(2)} \in L_{p,\psi^{1-p}}(\mathbf{R})$ , satisfying

$$T_1^{(2)}f(y) = H_1^{(2)}(y),$$

for any  $y \in \mathbf{R}$ .

In view of (5.9), it follows that  $\|T_1^{(2)}f\|_{p,\psi^{1-p}} = \|H_1^{(2)}\|_{p,\psi^{1-p}} \leq K_1(\sigma) \|f\|_{p,\phi}$ , and then the operator  $T_1^{(2)}$  is bounded satisfying

$$\|T_1^{(2)}\| = \sup_{f(\neq 0) \in L_{p,\phi}(\mathbf{R}_+)} \frac{\|T_1^{(2)}f\|_{p,\psi^{1-p}}}{\|f\|_{p,\phi}} \leq K_1(\sigma).$$

Since the constant factor  $K_1(\sigma)$  in (5.9) is the best possible, we have  $\|T_1^{(2)}\| = K_1(\sigma)$ .

If we define the formal inner product of  $T_1^{(2)}f$  and  $g$  as follows:

$$(T_1^{(2)}f, g) := \int_{-\infty}^{\infty} \left[ \int_{\tilde{E}_y} \frac{(\min\{|x|, |y|\})^\beta}{|x+y|^{\lambda+\beta}} f(x) dx \right] g(y) dy,$$

then we can rewrite (3.17) and (3.18) as follows:

$$(T_1^{(2)}f, g) < \|T_1^{(2)}\| \cdot \|f\|_{p,\phi} \|g\|_{q,\psi}, \|T_1^{(2)}f\|_{p,\psi^{1-p}} < \|T_1^{(2)}\| \cdot \|f\|_{p,\phi}. \quad (5.10)$$

(f) In view of Corollary 4.4, for  $f \in L_{p,\phi}(\mathbf{R})$ , setting

$$H_2^{(2)}(y) := \int_{-|y|}^{|y|} \frac{(\min\{|x|, |y|\})^\beta}{|x+y|^{\lambda+\beta}} |f(x)| dx \quad (y \in \mathbf{R}),$$

by (4.1), we have

$$\|H_2^{(2)}\|_{p,\Psi^{1-p}} = \left[ \int_{-|y|}^{|y|} \Psi^{1-p}(y) (H_2^{(2)}(y))^p dy \right]^{\frac{1}{p}} < K_2(\sigma) \|f\|_{p,\phi} < \infty. \quad (5.11)$$

**Definition 5.6.** Define the Hilbert-type integral operator of the second kind with the homogeneous kernel in the whole plane  $T_2^{(2)} : L_{p,\phi}(\mathbf{R}) \rightarrow L_{p,\Psi^{1-p}}(\mathbf{R})$  as follows: For any  $f \in L_{p,\phi}(\mathbf{R})$ , there exists a unique representation  $T_2^{(2)} f = H_2^{(2)} \in L_{p,\Psi^{1-p}}(\mathbf{R})$ , satisfying

$$T_2^{(2)} f(y) = H_2^{(2)}(y),$$

for any  $y \in \mathbf{R}$ .

In view of (5.11), it follows that  $\|T_2^{(2)} f\|_{p,\Psi^{1-p}} = \|H_2^{(2)}\|_{p,\Psi^{1-p}} \leq K_2(\sigma) \|f\|_{p,\phi}$ , and thus the operator  $T_2^{(2)}$  is bounded satisfying

$$\|T_2^{(2)}\| = \sup_{f(\neq 0) \in L_{p,\phi}(\mathbf{R}_+)} \frac{\|T_2^{(2)} f\|_{p,\Psi^{1-p}}}{\|f\|_{p,\phi}} \leq K_2(\sigma).$$

Since the constant factor  $K_2(\sigma)$  in (5.11) is the best possible, we have  $\|T_2^{(2)}\| = K_2(\sigma)$ .

If we define the formal inner product of  $T_2^{(2)} f$  and  $g$  as

$$(T_2^{(2)} f, g) := \int_{-\infty}^{\infty} \left[ \int_{-|y|}^{|y|} \frac{(\min\{|x|, |y|\})^\beta}{|x+y|^{\lambda+\beta}} f(x) dx \right] g(y) dy,$$

then we can rewrite (3.19) and (4.1) as follows:

$$(T_2^{(2)} f, g) < \|T_2^{(2)}\| \cdot \|f\|_{p,\phi} \|g\|_{q,\Psi} \|T_2^{(2)} f\|_{p,\Psi^{1-p}} < \|T_2^{(2)}\| \cdot \|f\|_{p,\phi}. \quad (5.12)$$

## Acknowledgments

The authors wish to express their thanks to Professors Mario Krnic and Jichang Kuang for their careful reading of the manuscript and their valuable suggestions.

## References

- [1] G.H. Hardy, J.E. Littlewood, G.Pólya, *Inequalities*, Cambridge University Press, Cambridge, USA, 1934.
- [2] D.S. Mitrinović, J.E. Pecaric, A.M.Fink, *Inequalities Involving Functions and their Integrals and Derivatives*, Kluwer Academic, Boston, USA, 1991.

- [3] B.C. Yang, The Norm of Operator and Hilbert-Type Inequalities, Science Press, Beijing, China, 2009.
- [4] B.C. Yang, A survey of the study of Hilbert-type inequalities with parameters, *Advances in Mathematics*, 38(3) (2009), 257-268.
- [5] B.C. Yang, On the norm of an integral operator and applications, *J. Math. Anal. Appl.*, 321(2006), 182-192.
- [6] J.S. Xu, Hardy-Hilbert's inequalities with two parameters, *Advances in Mathematics*, 36(2)(2007), 63-76.
- [7] B.C. Yang, On the norm of a Hilbert's type linear operator and applications, *J. Math. Anal. Appl.*, 325(2007), 529-541.
- [8] D.M. Xin, A Hilbert-type integral inequality with the homogeneous kernel of zero degree, *Mathematical Theory and Applications*, 30(2)(2010), 70-74.
- [9] B.C. Yang, A Hilbert-type integral inequality with the homogenous kernel of degree 0, *Journal of Shandong University (Natural Science)*, 45(2)(2010), 103-106.
- [10] L. Debnath, B.C. Yang, Recent developments of Hilbert-type discrete and integral inequalities with applications, *International Journal of Mathematics and Mathematical Sciences*, Volume 2012, Article ID 871845, 29 pages.
- [11] G.V. Milovanović, M.Th. Rassias, Some properties of a hypergeometric function which appear in an approximation problem, *Journal of Global Optimization*, 57(2013), 1173-1192.
- [12] M.Th. Rassias, B.C. Yang, On half-discrete Hilbert's inequality. *Applied Mathematics and Computation*, 220(2.13), 75 - 93.
- [13] Th.M. Rassias, B.C. Yang, A multidimensional half - discrete Hilbert - type inequality and the Riemann zeta function, *Applied Mathematics and Computation*, 225(2013), 263-277.
- [14] M.Th. Rassias, B.C. Yang, On a multidimensional half - discrete Hilbert - type inequality related to the hyperbolic cotangent function. *Applied Mathematics and Computation*, 242(2013), 800-813.
- [15] M.Th. Rassias, B.C. Yang, A multidimensional Hilbert - type integral inequality related to the Riemann zeta function, *Applications of Mathematics and Informatics in Science and Engineering* (N. J. Daras, ed.), Springer, New York, 417-433, 2014.
- [16] G.V. Milovanović, M.Th. Rassias (eds.), *Analytic Number Theory, Approximation Theory and Special Functions*, Springer, New York, 2014.
- [17] B.C. Yang, A new Hilbert-type integral inequality, *Soochow Journal of Mathematics*, 33(4)(2007), 849-859.

- [18] B.C. Yang, A Hilbert-type integral inequality with a non-homogeneous kernel, *Journal of Xiamen University (Natural Science)*, 48(2)(2008), 165-169.
- [19] B. He, B.C. Yang, On a Hilbert-type integral inequality with the homogeneous kernel of 0-degree and the hypergeometric function, *Mathematics in Practice and Theory*, 40(18)(2010), 203-211.
- [20] B.C. Yang, A new Hilbert-type integral inequality with some parameters, *Journal of Jilin University (Science Edition)*, 46(6)(2008), 1085-1090.
- [21] Z. Zeng, Z.T Xie, On a new Hilbert-type integral inequality with the homogeneous kernel of degree 0 and the integral in whole plane, *Journal of Inequalities and Applications*, Vol. 2010, Article ID 256796, 9 pages.
- [22] B.C. Yang, A reverse Hilbert-type integral inequality with some parameters, *Journal of Xinxiang University (Natural Science Edition)*, 27(6)(2010), 1-4.
- [23] A.Z. Wang, B.C. Yang, A new Hilbert-type integral inequality in whole plane with the non-homogeneous kernel, *Journal of Inequalities and Applications*, Vol. 2011, 2011: 123, doi:10.1186/1029-24X-2011-123.
- [24] D.M. Xin, B.C. Yang, A Hilbert-type integral inequality in whole plane with the homogeneous kernel of degree -2, *Journal of Inequalities and Applications*, Vol. 2011, Article ID 401428, 11 pages.
- [25] B. He, B.C. Yang, On an inequality concerning a non-homogeneous kernel and the hypergeometric function, *Tamsul Oxford Journal of Information and Mathematical Sciences*, 27(1)(2011), 75-88.
- [26] B.C. Yang, A reverse Hilbert-type integral inequality with a non-homogeneous kernel, *Journal of Jilin University (Science Edition)*, 49(3)(2011), 437-441.
- [27] Q.L. Huang, S.H. Wu, B.C. Yang, Parameterized Hilbert-type integral inequalities in the whole plane, *The Scientific World Journal*, Volume 2014, Article ID 169061, 8 pages.
- [28] Z. Zhen, K. Raja Rama Gandhi, Z.T. Xie, A new Hilbert-type inequality with the homogeneous kernel of degree -2 and with the integral, *Bulletin of Mathematical Sciences & Applications*, 3(1)(2014), 11-20.
- [29] Z.X. Wang, D.R. Guo, *Introduction to Special Functions*, Science Press, Beijing, China, 1979.
- [30] J.C. Kuang, *Applied Inequalities*. Shangdong Science and Technology Press, Jinan, China, 2004.
- [31] J.C. Kuang, *Introduction to Real Analysis*. Hunan Education Press, Changsha, China, 1996.